

SD-A088 181

SOUTH CAROLINA UNIV COLUMBIA DEPT OF MATHEMATICS COM--ETC F/6 12/1
THE ANALYSIS OF A RANDOMIZED COMPLETE BLOCK DESIGN WHEN OBSERVA--ETC(U)
NOV 79 L J WEI F49620-79-C-0140

UNCLASSIFIED

AFOSR-TR-80-0568

NL

1 of 1
95.3
10/10/80



END
DATE
FILMED
9-80
DTIC

18

19

AFOSR-TR-80-0568

5

AD A088181

LEVEL #

2. Introduction

6

THE ANALYSIS OF A RANDOMIZED COMPLETE BLOCK DESIGN WHEN OBSERVATIONS ARE SUBJECT TO ARBITRARY RIGHT CENSORSHIP.

10 L. J. Wei

University of South Carolina

11 Nov 1979

12 17

DTIC ELECTE
AUG 19 1980

A

University of South Carolina
Department of Mathematics,
Computer Science, and Statistics
Columbia, South Carolina 29208

16 2344

17 A5

DDC FILE COPY

*Research supported by the United States Air Force Office of Scientific Research under Contract No. F49620-79-C-0140

16

80 8 14 118

Approved for public release;
distribution unlimited.


449442

THE ANALYSIS OF A RANDOMIZED COMPLETE BLOCK DESIGN
WHEN OBSERVATIONS ARE SUBJECT TO ARBITRARY RIGHT CENSORSHIP

L. J. Wei

Department of Mathematics, Computer Science, and Statistics
University of South Carolina

Based on arbitrarily right censored observations, asymptotically distribution free tests are proposed for testing the equivalence of K treatments in the analysis of a randomized complete block design with random block effect. Multiple comparisons procedures which incorporate a pairwise ranking scheme are also proposed for determining which, if any, treatments differ from each other.



Key words and phrases: Asymptotically distribution free test; Multiple comparisons procedure; Friedman test; Ordered alternative; Gehan statistic.

AIR FORCE OFFICE OF SCIENTIFIC RESEARCH (AFSC)
NOTICE OF TRANSMITTAL TO DDC
This report has been reviewed and is
approved for distribution in accordance with AFM 190-12 (7b).
Distribution is unlimited.
A. D. BLOSS
Technical Information Officer

1. INTRODUCTION

Let $(x_{11}^0, \dots, x_{k1}^0)', \dots, (x_{1n}^0, \dots, x_{kn}^0)'$ be independent, identically distributed random vectors with a continuous joint distribution function G^0 and marginal distribution functions F_i^0 , $i = 1, \dots, K$. The null hypothesis H_0 , which we wish to test, is that G^0 is symmetric in its K arguments, i.e. that

$$G^0(x_{i_1}^0, \dots, x_{i_K}^0) = G^0(x_1^0, \dots, x_K^0),$$

for all permutations (i_1, \dots, i_K) of $(1, \dots, K)$. Note that H_0 implies that

$$F_1^0(x) = F_2^0(x) = \dots = F_K^0(x), \text{ for all } x \in R.$$

Such a hypothesis testing problem arises when we test the equivalence of K treatments in the analysis of a randomized complete block design and assume that the blocks themselves are drawn at random from a population of blocks. There are several standard nonparametric tests for this case, for example, Friedman test and the aligned rank order test (Mehra & Sarangi (1967), and Sen (1968)). For an ordered alternative H_1 :

$$F_1^0(x) \leq F_2^0(x) \leq \dots \leq F_K^0(x), \text{ for all } x \in R,$$

where at least one of the inequalities is strict for some x , we have tests proposed by Doksum (1967) and Hollander (1967). If one is not only interested in a single global test of whether the K treatments are equivalent, then multiple comparisons procedures are more appropriate (cf Miller (1966, 1977)).

Now suppose that a clinical trial is conducted to compare K treatments, the parameter of interest is the length of survival. It is common that at

the end of the trial, there may be incomplete survival information on certain individuals. More specifically, the response X_{ij}^0 may be censored from right by an independent variable Z_{ij} so that the n K -tuples $(X_{ij}^0, \dots, X_{Kj}^0)'$ cannot always be observed. Rather our observations consist of nK pairs $(X_{ij}, \Delta_{ij})'$, where $X_{ij} = \min(X_{ij}^0, Z_{ij})$ and $\Delta_{ij} = 1$, if X_{ij}^0 is observed and zero otherwise. Furthermore, for mathematical convenience Z_{ij} is assumed to be governed by a continuous distribution function $J_i(\cdot)$, $i = 1, \dots, K$, $j = 1, \dots, n$.

In the parametric statistics, the procedure for testing H_0 based on observations $(X_{ij}, \Delta_{ij})'$ is complicated and no results have been obtained. Although a generalized Friedman test (Patel (1975)) is available for testing H_0 based on censored observations, it is rather inefficient when there are too many censored K -tuples in the data. As an extreme example, for the data $(12^+, 9^+, 13)'$, $(9, 8^+, 5^+)'$, $(4, 3^+, 2^+)'$, $(2, 1^+, 1^+)'$, where "+" denotes censoring, the generalized Friedman test which uses information only within blocks leads to no conclusion about H_0 .

In this article, we propose and analyze test procedures which allow interblock comparisons for testing H_0 based on observations $(X_{ij}, \Delta_{ij})'$. A multiple comparisons procedure which selects treatments, if any, that differ from one another in terms of enhanced survival is also presented.

The corresponding K sample problems have been investigated by Breslow (1970), Crowley & Thomas (1975), and Koziol & Reid (1977).

2. DEFINITIONS AND A BASIC THEOREM

Let \bar{F}_i^e and \bar{F}_i be the empirical and the theoretical survival functions of X_{ij} . Also, let \tilde{F}_i^e and \tilde{F}_i be the empirical and the theoretical subdistribution functions of uncensored X_{ij} , i.e.

$$\tilde{F}_i(s) = P(X_{ij} \leq s, \Delta_{ij} = 1) \quad \text{and}$$

$$\tilde{F}_i^e(s) = n^{-1} \sum_{j=1}^n I(X_{ij} \leq s, \Delta_{ij} = 1),$$

where $I(A)$ is the indicator function of the event A , $i = 1, \dots, K$.

Furthermore, the joint subdistribution functions of $(X_{ij}, X_{i',j})'$, $(X_{ij}, X_{i',j}, X_{kj}, X_{k',j})'$ and $(X_{ij}, \dots, X_{Kj})'$ are defined by

$$\tilde{D}_{ii'}(s, t; \delta, \epsilon) = P(X_{ij} \leq s, X_{i',j} \leq t, \Delta_{ij} = \delta, \Delta_{i',j} = \epsilon),$$

$$\tilde{H}_{ii'kk'}(s, t, s', t'; \delta, \epsilon, \delta', \epsilon') = P(X_{ij} \leq s, X_{i',j} \leq t, X_{kj} \leq s', X_{k',j} \leq t',$$

$$\Delta_{ij} = \delta, \Delta_{i',j} = \epsilon, \Delta_{kj} = \delta', \Delta_{k',j} = \epsilon'), \quad \text{and}$$

$$\tilde{G}(s_1, \dots, s_K; \delta_1, \dots, \delta_K) = P(X_{1j} \leq s_1, \dots, X_{Kj} \leq s_K, \Delta_{1j} = \delta_1, \dots, \Delta_{Kj} = \delta_K),$$

respectively,

where i, i', k , and k' denote distinct indices. The corresponding empirical subdistribution functions are defined by

$$\tilde{D}_{ii'}^e(s, t; \delta, \epsilon) = n^{-1} \sum_{j=1}^n I(X_{ij} \leq s, X_{i',j} \leq t, \Delta_{ij} = \delta, \Delta_{i',j} = \epsilon),$$

$$\tilde{H}_{ii'kk'}^e(s, t, s', t'; \delta, \epsilon, \delta', \epsilon') =$$

$$n^{-1} \sum_{j=1}^n I(X_{ij} \leq s, X_{i',j} \leq t, X_{kj} \leq s', X_{k',j} \leq t', \Delta_{ij} = \delta, \Delta_{i',j} = \epsilon, \Delta_{kj} = \delta', \Delta_{k',j} = \epsilon')$$

and

$$\tilde{G}^e(s_1, \dots, s_K; \delta_1, \dots, \delta_K) = n^{-1} \sum_{j=1}^n I(X_{1j} \leq s_1, \dots, X_{Kj} \leq s_K, \Delta_{1j} = \delta_1, \dots, \Delta_{Kj} = \delta_K).$$

All the procedures proposed in this article are based on the Gehan scoring function ψ , where, for any two pairs of observations $(X_{ij}, \Delta_{ij})'$ and $(X_{i'j'}, \Delta_{i'j'})'$,

$$\psi(X_{ij}, \Delta_{ij}, X_{i'j'}, \Delta_{i'j'}) = \begin{cases} +1, & X_{ij} > X_{i'j'}, \Delta_{i'j'} = 1 \\ -1, & X_{ij} < X_{i'j'}, \Delta_{ij} = 1 \\ 0, & \text{otherwise.} \end{cases}$$

Now, let $p_{ii'} = E\psi(X_{ik}, \Delta_{ik}, X_{i'k'}, \Delta_{i'k'})$, where $i \neq i'$ and $k \neq k'$.

Under H_0 , $p_{ii'} = 0$ which is independent of censoring distributions J_i and $J_{i'}$ (cf Efron (1967) and Mantel (1967)). Consider Gehan's statistic (1965) $V_{ii',n}$ for populations i and i' :

$$V_{ii',n} = n^{-\frac{3}{2}} \sum_{j=1}^n \sum_{j'=1}^n \psi(X_{ij}, \Delta_{ij}, X_{i'j'}, \Delta_{i'j'}).$$

The following Theorem whose proof is in the Appendix provides the asymptotic

distribution of random vectors $\langle V_{ii,n}^* \rangle$, where $V_{ii,n}^* = n^{-\frac{3}{2}} \sum_{j=1}^n \sum_{j'=1}^n (\psi(X_{ij}, \Delta_{ij}, X_{i'j'}, \Delta_{i'j'}) - p_{ii'})$, $i, i' = 1, \dots, K$ and $i < i'$.

Theorem 1. As $n \rightarrow \infty$, the random vector $\langle V_{ii,n}^* \rangle$ converges in distribution to a multivariate normal random vector with mean 0 and covariance matrix $\Sigma = (\sigma_{ii',kk'})$, $i < i', k < k'$. Under H_0 ,

$$\sigma_{ii',kk'} = \sigma_{ii',kk'}^0 =$$

$$E\left(\tilde{F}_1(X_{1j}) - \Delta_{1j}\bar{F}_1(X_{1j}) + \Delta_{1,j}\bar{F}_1(X_{1,j}) - \tilde{F}_1(X_{1,j})\right) \\ \left(\tilde{F}_k(X_{kj}) - \Delta_{kj}\bar{F}_k(X_{kj}) + \Delta_{k,j}\bar{F}_k(X_{k,j}) - \tilde{F}_k(X_{k,j})\right) .$$

If the censoring variables Z_{1j}, \dots, Z_{Kj} are interchangeable (hereafter referred to as Condition A), then $J_1 = J_2 = \dots = J_K$ and, under H_0 , the random pairs $(X_{1j}, \Delta_{1j})', \dots, (X_{Kj}, \Delta_{Kj})'$ are also interchangeable, $j = 1, \dots, n$. For future references, let $\tilde{F} = \tilde{F}_1 = \dots = \tilde{F}_K$ and $\bar{F} = \bar{F}_1 = \dots = \bar{F}_K$ and also let

$$\sigma^2 = E\left(\tilde{F}(X_{1j}) - \Delta_{1j}\bar{F}(X_{1j})\right)^2 \text{ and}$$

$$\tau = E\left(\tilde{F}(X_{1j}) - \Delta_{1j}\bar{F}(X_{1j})\right)\left(\tilde{F}(X_{1,j}) - \Delta_{1,j}\bar{F}(X_{1,j})\right) .$$

3. MULTIPLE COMPARISONS UNDER H_0

When it is less important to test the overall hypothesis H_0 than to compare certain populations on an individual basis, a multiple comparisons procedure is more desirable than a single global test. The decision of which, if any, populations differ from one another can be made by examining the individual comparison $V_{ii',n}$. We would conclude that populations i and i' differ if $|V_{ii',n}|$ is sufficiently large. A conservative multiple comparisons procedure at level α can be obtained through an inequality due to Šidák (1967), i.e. if $\underline{U} = (U_1, \dots, U_r)'$ is normally distributed with mean \underline{Q} and correlation matrix R , then

$$P_r \left(\max_{1 \leq i \leq r} |U_i| \leq u \mid U \sim N_r(Q, R) \right) \geq$$

$$P_r \left(\max_{1 \leq i \leq r} |U_i| \leq u \mid U \sim N_r(Q, I_r) \right) .$$

It follows from Theorem 1 that, for large n , one would judge that populations i and i' as different at level α if

$$|v_{ii',n}| / (\hat{\sigma}_{ii',ii'}^0)^{1/2} > \xi_{\beta/2} ,$$

where $\xi_{\beta/2}$ is the $100(1-\beta/2)$ percentage point of the $N(0,1)$ distribution, $(1-\beta)^{K(K-1)/2} = 1-\alpha$, and $\hat{\sigma}_{ii',ii'}^0$ is a (strongly) consistent estimator of $\sigma_{ii',ii'}^0$:

$$\hat{\sigma}_{ii',ii'}^0 = \sum_{\delta=0}^1 \sum_{\epsilon=0}^1 \int [\tilde{F}_1^e(s) - \delta \bar{F}_1^e(s) + \epsilon \bar{F}_1^e(t) - \tilde{F}_1^e(t)]^2 d\tilde{D}_{ii'}^e(s,t;\delta,\epsilon) .$$

Another multiple comparisons procedure can be obtained through the Bonferroni inequality. But as indicated by Koziol and Reid (1977, p. 1154), this procedure is less powerful than the one based on Šidák inequality.

Some improvement of the above results can be made under Condition A (the interchangeability of Z_{1j}, \dots, Z_{Kj}). It can be shown that, under H_0 and Condition A, the asymptotic covariance structure of $\langle \frac{v_{ii',n}}{(\sigma_{ii',ii'}^2 - \tau)^{1/2}} \rangle$ is identical to that of the vector having $K(K-1)/2$ components $Y_i - Y_{i'}$, $1 \leq i < i' \leq K$, where $Y = (Y_1, \dots, Y_K)' \sim N_K(Q, I_K)$, and I_K is the identity matrix. Therefore, one would conclude that populations i and i' differ at level α if

$$\frac{|v_{ii',n}|}{(\hat{\sigma}_{ii',ii'}^2 - \hat{\tau}_n)^{1/2}} > q_K^\alpha ,$$

where q_K^α is the $100(1-\alpha)$ percentage point of the distribution of the range of K independent unit normal random variables (cf. Pearson and Hartley (1966), Tables 22 and 23) and $\hat{\sigma}_n^2$ and $\hat{\tau}_n$ are (strongly) consistent estimators of σ^2 and τ :

$$\begin{aligned}\hat{\sigma}_n^2 &= \int (\tilde{F}^e(s) - \bar{F}^e(s))^2 d\tilde{F}^e(s) - \int (\tilde{F}^e(s))^2 d(\bar{F}^e(s) + \tilde{F}^e(s)) , \\ \binom{K}{2} \hat{\tau}_n &= \sum_{\delta, \epsilon=0}^1 \sum_{i=1}^{K-1} \sum_{i'=i+1}^K \int (\tilde{F}^e(s) - \delta \bar{F}^e(s)) (\tilde{F}^e(t) - \epsilon \bar{F}^e(t)) d\tilde{D}_{ii'}^e(s, t; \delta, \epsilon) , \\ \tilde{F}^e(s) &= K^{-1} \sum_{i=1}^K \tilde{F}_i^e(s) \quad \text{and} \quad \bar{F}^e(s) = K^{-1} \sum_{i=1}^K \bar{F}_i^e(s) .\end{aligned}$$

4. TESTING H_0 AGAINST ORDERED ALTERNATIVE H_1

In this section, we present a test statistic S_n which is a generalization of the Jonckheere K -sample test (Jonckheere (1954), Patel & Hoel (1973)) for testing H_0 against H_1 based on observations $(X_{ij}, \Delta_{ij})'$, where

$$S_n = \sum_{i=1}^{K-1} \sum_{i'=i+1}^K v_{ii',n} .$$

Large values of S_n suggest rejection of the null hypothesis H_0 . It follows from Theorem 1 that, under H_0 , S_n converges in distribution to a normal random variable with mean 0 and variance

$$\sum_{i=1}^{K-1} \sum_{i'=i+1}^K \sum_{k=1}^{K-1} \sum_{k'=k+1}^K \sigma_{ii',kk'}^0 .$$

Note that $\sigma_{ii',kk'}^0$, where i, i', k and k' are distinct integers, can be estimated consistently by

$$\hat{\sigma}_{ii',kk'}^0 = \sum_{\delta, \epsilon, \delta', \epsilon'=0}^1 \int \left(\tilde{F}_{i'}^e(s) - \delta \bar{F}_{i'}^e(s) + \epsilon \bar{F}_{i'}^e(t) - \tilde{F}_{i'}^e(t) \right) \left(\tilde{F}_k^e(s') - \delta' \bar{F}_k^e(s') + \epsilon' \bar{F}_k^e(t') - \tilde{F}_k^e(t') \right) d\tilde{H}_{ii',kk'}^e(s, t, s', t'; \delta, \epsilon, \delta', \epsilon') .$$

5. A GLOBAL TEST FOR H_0

If we are interested in testing H_0 against a general alternative hypothesis that G^0 is not symmetric in its K arguments, then a global test is more appropriate. Unfortunately, it is not feasible to get such a test under unequal patterns of censorship operative on K populations. In this Section, we assume that the censoring variables Z_{1j}, \dots, Z_{Kj} are interchangeable (Condition A) and present a generalized Breslow test (1970) for testing H_0 based on observations $(X_{ij}, \Delta_{ij})'$.

Define a vector score statistic $\tilde{W}_n = (W_{1n}, \dots, W_{Kn})'$, where W_{in} is the total score comparing the i th sample with the remaining $(K-1)$ samples, i.e.

$$W_{in} = \sum_{\substack{i'=1 \\ i' \neq i}}^K v_{ii',n} .$$

It follows from Theorem 1 that, under H_0 , \tilde{W}_n converges in distribution to a normal random vector with mean $\underline{0}$ and covariance matrix

$$\Gamma = \begin{pmatrix} \gamma^2 & \eta & \dots & \eta \\ \eta & \gamma^2 & \dots & \eta \\ . & . & \dots & . \\ \eta & \eta & \dots & \gamma^2 \end{pmatrix}_{K \times K},$$

where $\gamma^2 = E\phi^2$, $\eta = E\phi\phi'$,

$$\phi = (K-1)(\tilde{F}(x_{1j}) - \Delta_{1j}\bar{F}(x_{1j})) + \sum_{m=2}^K \Delta_{mj}\bar{F}(x_{mj}) \quad \text{and}$$

$$\phi' = (K-1)(\tilde{F}(x_{Kj}) - \Delta_{Kj}\bar{F}(x_{Kj})) + \sum_{\ell=1}^{K-1} \Delta_{\ell j}\bar{F}(x_{\ell j}) .$$

Again, γ^2 and η can be estimated consistently, for example, by $\hat{\gamma}_n^2$ and $\hat{\eta}_n$, where

$$\hat{\gamma}_n^2 = \sum_{\delta_1, \dots, \delta_K=0} [(K-1)(\tilde{F}^e(s_1) - \delta_1 \bar{F}^e(s_1)) + \sum_{m=2}^K \delta_m \bar{F}^e(s_m)]^2 d\tilde{G}^e(s_1, \dots, s_K; \delta_1, \dots, \delta_K) ,$$

and

$$\begin{aligned} \hat{\eta}_n = \sum_{\delta_1, \dots, \delta_K=0} \int & [(K-1)(\tilde{F}^e(s_1) - \delta_1 \bar{F}^e(s_1)) + \sum_{m=2}^K \delta_m \bar{F}^e(s_m)] \\ & [(K-1)(\tilde{F}^e(s_K) - \delta_K \bar{F}^e(s_K)) + \sum_{\ell=1}^{K-1} \delta_\ell \bar{F}^e(s_\ell)] d\tilde{G}^e(s_1, \dots, s_K; \delta_1, \dots, \delta_K) . \end{aligned}$$

The latent roots of Γ are $(\gamma^2 + (K-1)\eta)$, which is zero, with the latent vector $q_K = (1, \dots, 1)' / K^{1/2}$ and $(\gamma^2 - \eta)$ of multiplicity $(K-1)$ with $(K-1)$ latent (any set of) vectors orthogonal to $(1, \dots, 1)'$, q_1, \dots, q_{K-1} , say (cf Rao (1973), p. 67). For example, when $K = 3$, $q_1 = (\frac{3}{2})^{1/2}(-\frac{1}{2}, 1, -\frac{1}{2})'$, $q_2 = (\frac{1}{2})^{1/2}(\frac{1}{2}, -\frac{1}{2}, 0)'$, $q_3 = (\frac{1}{3})^{1/2}(1, 1, 1)'$. In general, the q_i are perhaps most easily found by means of the Gram-Schmidt orthogonalization process. Since Γ is symmetric, therefore, $Q'\Gamma Q = \Lambda_{K \times K}$, where $Q = (q_1, \dots, q_K)$ which is an orthogonal matrix, $\Lambda_{K \times K} = \text{diag}(\lambda, \dots, \lambda, 0)$ and $\lambda = \gamma^2 - \eta$. It follows that, under H_0 ,

$$T_n = \hat{\lambda}_n^{-1} \sum_{i=1}^{K-1} (q_i' W_n)^2$$

is asymptotically distributed in a chi-squared distribution with $K-1$ degrees

of freedom, where $\hat{\lambda}_n = \hat{\gamma}_n^2 - \hat{\eta}_n$. Large values of T_n suggest a rejection decision of H_0 .

For $K = 2$ and unequal patterns of censorship, an asymptotically distribution free test was studied by Wei (1979) for testing H_0 based on observations $(X_{ij}, \Delta_{ij})'$, $i = 1, 2$; $j = 1, \dots, n$. This test is reduced to the above generalized Breslow test when the two censoring distributions J_1 and J_2 are equal.

6. REMARKS

In practice, all the consistent estimators in this article can be obtained through simple summation processes. Therefore, it is feasible to evaluate the test procedures by hand calculations and extremely easy if a small programmable calculator is available.

Although there are several multivariate parametric models available for life testing in reliability (cf Freund (1961), Marshall & Olkin (1967), and Block & Basu (1974)), none is satisfactory for survivability theory, especially for our current settings. For example, there is no continuous multivariate distribution function which has exponential marginals and possesses a physical interpretation in survival analysis. This probably makes nonparametric procedures more appealing in multivariate survival studies.

Also, for lack of appropriate parametric models, the efficiency study of our new test procedures seems unfeasible. However, since the new procedures allow interblock comparisons, we expect that they are more powerful than the procedures based on Friedman statistic.

7. APPENDIX

Proof of Theorem 1.

Let

$$\psi_{ii'}^0(X_{ij}, \Delta_{ij}) = E[\psi(X_{ij}, \Delta_{ij}, X_{i'j'}, \Delta_{i'j'}) | X_{i'j'}, \Delta_{i'j'}] - P_{ii'} \quad \text{and}$$

$$\psi_{ii'}^1(X_{i'j'}, \Delta_{i'j'}) = E[\psi(X_{ij}, \Delta_{ij}, X_{i'j'}, \Delta_{i'j'}) | X_{ij}, \Delta_{ij}] - P_{ii'} \quad , \quad \text{where } j \neq j'.$$

Also, let

$$v_{ii',n}^{**} = n^{-1/2} \left(\sum_{j=1}^n (\psi_{ii'}^0(X_{ij}, \Delta_{ij}) + \psi_{ii'}^1(X_{i'j'}, \Delta_{i'j'})) \right)$$

and

$$g(X_{ij}, \Delta_{ij}, X_{i'j'}, \Delta_{i'j'}) = \psi(X_{ij}, \Delta_{ij}, X_{i'j'}, \Delta_{i'j'}) - P_{ii'} - \\ \psi_{ii'}^0(X_{ij}, \Delta_{ij}) - \psi_{ii'}^1(X_{i'j'}, \Delta_{i'j'}) .$$

Then,

$$E(v_{ii',n}^* - v_{ii',n}^{**})^2 = n^{-3} E \left[\sum_{j=1}^n \sum_{j'=1}^n g(X_{ij}, \Delta_{ij}, X_{i'j'}, \Delta_{i'j'}) \right]^2 \\ = n^{-3} \sum_{j=1}^n \sum_{j'=1}^n \sum_{\ell=1}^n \sum_{\ell'=1}^n h(j, j', \ell, \ell') , \quad (A.1)$$

where $h(j, j', \ell, \ell') = E[g(X_{ij}, \Delta_{ij}, X_{i'j'}, \Delta_{i'j'}) g(X_{i\ell}, \Delta_{i\ell}, X_{i'\ell'}, \Delta_{i'\ell'})]$.

Because g is bounded, we can ignore any u terms of the sum in (A.1) if u is of order $O(n^3)$. Consider the following cases for which the number of terms is with order larger than or equal to $O(n^3)$ (where j, j', ℓ and ℓ' represent four distinct indices):

$$(1) \quad h(j, j', \ell, \ell') = E g(X_{ij}, \Delta_{ij}, X_{i'j'}, \Delta_{i'j'}) E g(X_{i\ell}, \Delta_{i\ell}, X_{i'\ell'}, \Delta_{i'\ell'}) = 0 ;$$

$$(2) \quad h(j, j, \ell, \ell') = h(j, j', \ell, \ell) = 0 ;$$

$$(3) \quad h(j, j', j, \ell') = h(j, j', \ell, j')$$

$$\begin{aligned} &= E \{ E [g(X_{ij}, \Delta_{ij}, X_{i'j'}, \Delta_{i'j'}) g(X_{ij}, \Delta_{ij}, X_{i'\ell'}, \Delta_{i'\ell'}) | X_{ij}, \Delta_{ij}] \} \\ &= E \{ E [g(X_{ij}, \Delta_{ij}, X_{i'j'}, \Delta_{i'j'}) | X_{ij}, \Delta_{ij}] E [g(X_{ij}, \Delta_{ij}, X_{i'\ell'}, \Delta_{i'\ell'}) | X_{ij}, \Delta_{ij}] \} = 0 ; \end{aligned}$$

$$\begin{aligned} (4) \quad h(j, j', \ell, j) &= h(j, j', j', \ell') = E E [g(X_{ij}, \Delta_{ij}, X_{i'j'}, \Delta_{i'j'}) g(X_{i\ell}, \Delta_{i\ell}, X_{i'j'}, \Delta_{i'j'}) \\ &\quad | X_{ij}, \Delta_{ij}, X_{i'j'}, \Delta_{i'j'}] = E \{ E [g(X_{ij}, \Delta_{ij}, X_{i'j'}, \Delta_{i'j'}) | X_{ij}, \Delta_{ij}, X_{i'j'}, \Delta_{i'j'}] \\ &\quad E [g(X_{i\ell}, \Delta_{i\ell}, X_{i'j'}, \Delta_{i'j'}) | X_{ij}, \Delta_{ij}, X_{i'j'}, \Delta_{i'j'}] \} = 0 . \end{aligned}$$

It follows that $E(V_{ii',n}^* - V_{ii',n}^{**})^2 \rightarrow 0$, as $n \rightarrow \infty$. By Corollary 6 of Lehmann (1975), p. 289) and the fact that the vector $\langle V_{ii',n}^{**} \rangle$ has a multivariate normal limiting distribution $N(\underline{0}, \underline{I})$, the vector $\langle V_{ii',n}^* \rangle$ has the same multivariate normal limiting distribution $N(\underline{0}, \underline{I})$. Under H_0 , $p_{ii'} = 0$, it follows that $V_{ii',n} = V_{ii',n}^*$,

$$\psi_{ii'}^0(X_{ij}, \Delta_{ij}) = \tilde{F}_i(X_{ij}) - \Delta_{ij} \bar{F}_i(X_{ij}) \quad \text{and}$$

$$\psi_{ii'}^1(X_{i'j'}, \Delta_{i'j'}) = \Delta_{i'j'} \bar{F}_i(X_{i'j'}) - \tilde{F}_i(X_{i'j'}) .$$

Q.E.D.

REFERENCES

- Block, H. W. and Basu, A. P. (1974). A continuous bivariate exponential extension. J. Amer. Statist. Assoc. 69, 1031-1037.
- Breslow, N. (1970). A generalized Kruskal-Wallis test for comparing K samples subject to unequal patterns of censorship. Biometrika 57, 579-594.
- Crowley, J. and Thomas, D. R. (1975). Large sample theory for the log rank test. Tech. Report 415, Dept. of Statist., Univ. of Wisconsin, Madison, Wisconsin.
- Doksum, K. (1967). Robust procedures for some linear models with one observation per cell. Ann. Math. Statist. 38, 878-883.
- Efron, B. (1967). The two sample problem with censored data. Proc. Fifth Berkeley Symp. 4, 831-854.
- Freund, J. E. (1961). A bivariate extension of the exponential distribution. J. Amer. Statist. Assoc. 56, 971-977.
- Gehan, E. (1965). A generalized Wilcoxon test for comparing arbitrarily single censored samples. Biometrika 52, 203-223.
- Hollander, M. (1967). Rank tests for randomized blocks when the alternatives have an a priori ordering. Ann. Math. Statist. 38, 867-877.
- Jonckheere, A. R. (1954). A distribution free K-sample test against ordered alternative. Biometrika 41, 133-145.
- Koziol, J. A. and Reid, N. (1977). On multiple comparisons among K samples subject to unequal patterns of censorship. Commu. in Statist. A6(12), 1149-1164.
- Lehmann, E. L. (1975). Nonparametrics: Statistical Methods Based on Ranks. San Francisco, Cal.: Holden-Day, Inc.
- Mantel, N. (1967). Ranking procedures for arbitrarily restricted observations. Biometrics 23, 65-78.
- Marshall, A. W. and Olkin, I. (1967). A multivariate exponential distribution. J. Amer. Statist. Assoc. 62, 30-44.

- Mehra, K. L. and Sarangi, J. (1967). Asymptotic efficiency of certain rank tests for comparative experiments. Ann. Math. Statist. 38, 90-107.
- Miller, R. G., Jr. (1966). Simultaneous Statistical Inference, New York: McGraw-Hill Book Co.
- _____ (1977). Developments in multiple comparisons 1966-1976, J. of Amer. Statist. Assoc. 72, 779-788.
- Patel, K. M. and Hoel, D. G. (1973). A generalized Jonckheere K-sample test against ordered alternatives when observations are subject to arbitrary right censorship. Comm. in Statist. A2(4), 373-380.
- Patel, K. M. (1975). A generalized Friedman test for randomized block design when observations are subject to arbitrary right censorship. Comm. in Statist. A4(4), 389-395.
- Pearson, E. S. and Hartley, H. O. (1966). Biometrika Tables for Statisticians, Vol. I, 3rd Ed. Cambridge University Press.
- Rao, C. R. (1973). Linear Statistical Inference and its application, 2nd Ed., New York: John Wiley & Sons.
- Sen, P. K. (1968). On a class of aligned rank order tests in two way layouts. Ann. Math. Statist. 39, 1115-1124.
- Šidák (1967). Rectangular confidence regions for the means of multivariate normal distributions. J. Amer. Statist. Assoc. 62, 626-633.
- Wei, L. J. (1979). A generalized Gehan and Gilbert test for paired observations which are subject to arbitrary right censorship, Tech. Report, Univ. of South Carolina.

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER	2. GOVT ACCESSION NO. AD-A088 182	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) THE ANALYSIS OF A RANDOMIZED COMPLETE BLOCK DESIGN WHEN OBSERVATIONS ARE SUBJECT TO ARBITRARY RIGHT CONSORSHIP		5. TYPE OF REPORT & PERIOD COVERED Interim
7. AUTHOR(s) L. J. Wei		6. PERFORMING ORG. REPORT NUMBER
9. PERFORMING ORGANIZATION NAME AND ADDRESS University of South Carolina, Department of Mathematics, Computer Science, and Statistics ✓		8. CONTRACT OR GRANT NUMBER(s) F49620-79-C-0140 ✓
11. CONTROLLING OFFICE NAME AND ADDRESS Air Force Office of Scientific Research/NM Bolling AFB, Washington, D. C. 20332		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS 61102F 2034/A5
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		12. REPORT DATE November, 1979
		13. NUMBER OF PAGES 14
		15. SECURITY CLASS. (of this report) UNCLASSIFIED
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Asymptotically distribution free test; Multiple comparisons procedure; Friedman test; Ordered alternative; Gehan statistic.		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) Based on arbitrarily right censored observations, asymptotically distribution free tests are proposed for testing the equivalence of K treatments in the analysis of a randomized complete block design with random block effect. Multiple comparisons procedures which incorporate a pairwise ranking scheme are also proposed for determining which, if any, treatments differ from each other.		